

## Anti-h Transferrin 3105 SPTN-5

### Product overview

---

<b>Catalog number</b>	100440
<b>Specificity</b>	Antibody recognizes human transferrin
<b>Description</b>	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
<b>Product buffer solution</b>	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN <sub>3</sub> as a preservative
<b>Shelf life and storage</b>	36 months from manufacturing at 2–8 °C
<b>Subclass</b>	IgG <sub>1</sub>
<b>Analyte description</b>	Transferrins are glycoproteins which regulate the amount of free iron in circulation. Each transferrin molecule binds two Fe <sup>+3</sup> with high affinity. When no iron is bound the molecule is called Apotransferrin. Transferrin concentration elevates in anemia, but also during pregnancy and when using birth control pills. Chronic infections, malignant growth, alcohol induced liver damage, and other conditions with protein loss may lower transferrin concentration.

### Parameters tested on each lot

---

<b>Product appearance</b>	Liquid, may turn slightly opaque during storage
<b>Product concentration</b>	5.0 mg/ml (+/- 10%)
<b>Immunoreactivity</b>	80–120% compared to the reference sample in an FIA test
<b>IEF Profile</b>	6.5–7.6
<b>Purity</b>	≥ 95 %

### Kinetic parameters

---

<b>Association rate constant</b>	$6.7 \times 10^6$ 1/Ms
<b>Dissociation rate constant</b>	$1.3 \times 10^{-5}$ 1/s
<b>Affinity constant</b>	$K_A = 5.3 \times 10^{11}$ 1/M; $K_D = 1.9 \times 10^{-12}$ M (=0.002 nM)
<b>Determination method</b>	SPR analysis (ProteOn XPR36)
<b>Determination antigen</b>	Transferrin native human, AbD Serotec (Cat PHP-238, Lot 171210)



#### Legal disclaimer

**Cross-reactivities** Does not recognize lactoferrin or hemoglobin.

**Epitope** N/D

**Pair recommendations**

		DETECTION		
		3104	3105	3106
CAPTURE	3104	-	+	+
	3105	+	-	+
	3106	+	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

**Platforms tested** FIA

**Antigens tested** Apotransferrin (Sigma, T2252)

Product stability	TEMPERATURE, TIME	RESULT
	-70 °C, 21 days	OK
	-20 °C, 21 days	OK
	+4 °C, 21 days	OK
	+35 °C, 21 days	OK
	+45 °C, 3 days	OK
	+45 °C, 7 days	Minor charge alterations

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

**Miscellaneous** -

**References** -



**Legal disclaimer**