

Anti-h LH 5301 SP-5

Product overview

Catalog number	100016
Specificity	Antibody recognizes human luteinizing hormone (lutropin), and its beta-subunit
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
Product buffer solution	0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	36 months from manufacturing at 2–8 °C
Subclass	IgG ₁
Analyte description	In both males and females, LH is essential for reproduction. In females FSH initiates follicular growth and at the time of the maturation of the follicle the estrogen rise leads to a release of LH over a 24–48 hour period. This 'LH surge' triggers ovulation thereby not only releasing the egg, but also initiating the conversion of the residual follicle into a corpus luteum that, in turn, produces progesterone to prepare the endometrium for a possible implantation. LH is necessary to maintain luteal function for the first two weeks. In case of a pregnancy luteal function will be further maintained by the action of hCG from the newly established pregnancy. In the male, LH acts upon the Leydig cells of the testis and is responsible for the production of testosterone.

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/-10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	6.3–7.2
Purity	≥ 95 %

Kinetic parameters

Association rate constant	5.8×10^6 1/Ms
Dissociation rate constant	1.3×10^{-5} 1/s
Affinity constant	$K_A = 4.4 \times 10^{11}$ 1/M; $K_D = 2.3 \times 10^{-12}$ M (= 2.3 pM)
Determination method	SPR analysis (ProteOn XPR36)
Determination antigen	LH, Scripps Laboratories (Cat L0815)



Legal disclaimer

Cross-reactivities

LH α 13 % (Scripps Laboratories, Cat L0914, Lot 698811)
 LH β 170 % (Scripps Laboratories, Cat L1014, Lot 237711)
 FSH 5 % (Scripps Laboratories, Cat F0614, Lot 805811)
 hCG 138 % (Scripps Laboratories, Cat C0714, Lot 210164)
 TSH 0.03 % (Scripps Laboratories, Cat T0114, Lot 181711)

Epitope

Beta 2 as described in Nilsson et al. (2001)
 Two antibodies binding to the same, or closely located epitopes, belong to the same group and hence cannot be used as a pair in a sandwich assay.
 Epitope group numbering does not give any detailed information where the epitope is located.

Pair recommendations

		DETECTION				
		5301	5302	5303	5304	5501 (α subunit)
CAPTURE	5301	-	+	+	+	+
	5302	+	-	-	-	+
	5303	+	-	-	-	-
	5304	+	-	-	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA, CLIA, LF

Antigens tested Native LH antigen, Medix Biochemica 996-31.

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	Not Determined (N/D)
-20 °C, 21 days	N/D
+4 °C, 21 days	N/D
+35 °C, 7 days	N/D
+35 °C, 21 days	N/D
+45 °C, 3 days	N/D
+45 °C, 7 days	N/D

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous

Nilsson et al. (2001) analyzed epitopes of 30 different LH mAbs. Antibody 5301 was classified as belonging to epitope group beta 2, recognizing intact LH, its beta subunit as well as a common variant of LH. Antibody 5301 did not cross react with TSH or FSH but did so with hCG.



References

Federici, M.M., Fraser, R., Lundqvist, C., and Lankford, J.C., (1982) Production and characterization of monoclonal antibodies human lutenizing hormones. Fed. Proc., 41

Nilsson, C., Seppälä, M., and Pettersson, K., (2001) Immunological characterization of human luteinizing hormone with special regard to a common genetic variant. J.Endocrinol. 168:10-116

Pettersson, K.S.I., and Söderholm J.R-M., (1990) Ultrasensitive two-site immunometric assay of human lutropin by time-resolved fluorometry. Clin. Chem. 36(11):1928-1933

Pettersson, K.S.I., and Söderholm J.R-M., (1991) Individual differences in lutropin immunoreactivity revealed by monoclonal antibodies. Clin. Chem. 37(3):333-340

Vilja, P., Wichmann, L., Isola, J., and Tuohimaa, P., (1988) Monoclonal-antibody based noncompetitive avidin-biotin assay for lutropin in urine. Clin. Chem. 34(8):1585-1590



Legal disclaimer