


Recombinant Pepsinogen II, 100 µg

Catalog number	610001
Description	Recombinant human pepsinogen II protein. Predicted molecular weight: 41 kDa.
Amino acid sequence	MAVVKVPLKFKSIRETMKEKGLLGEFLRTHKYDPAWKYRFGDLSVTYEP MAYMDAAYFGEISIGTPPQNFLVLFDTGSSNLWVPSVYCQSQACTSHSRF NPSESSTYSTNGQTFSLQYSGSLTGFFGYDTLTVQSIQVPNQEFGLSEN EPGTNFVYAQFDGIMGLAYPALSVDEATTAMQGMVQEGALTSPVFSVYL SNQQGSSGGAVVFGGVDSSLYTGQIYWAPVTQELYWQIGIEEFLIGGQAS GWCSEGCQAIVDTGTSLLTVPQQYMSALLQATGAQEDEYGGFLVNCNSI QNLPSLTFIINGVEFPLPPSSYILSNNGYCTVGVPEPTYLSSQNGQPLWILGD VFLRSYYSVYDLGNRRVGFATAA
Product host	<i>Escherichia coli (E. coli)</i>
Product formulation	Lyophilized
Product buffer solution	Lyophilized from 50 mM Tris-HCl, pH 7.5; 150 mM NaCl, 0.5 µg/ml pepstatin A; containing 6 % sucrose as a stabilizer
Reconstitution	Reconstitute the lyophilized protein with deionized water
Recertification	36 months
Storage	Storage at 2–8 °C. Shelf life after reconstitution 3 months at 2–8 °C and 12 months at -20 °C.
Analyte description	Pepsinogen is the pro-form of pepsin and is produced in the stomach by chief cells. The major part of pepsinogen is secreted into the gastric lumen but a small amount can be found in the blood. Alterations in the serum pepsinogen concentrations has been found with <i>Helicobacter pylori (H. pylori)</i> infections, peptic ulcer disease, gastritis, and gastric cancer. More precise analysis may be achieved by measuring the pepsinogen I/II ratio.
Product concentration	N/A
Purity	Capillary electrophoresis (CE-SDS)



Antibodies tested	Anti-h Pepsinogen II 8101: + Anti-h Pepsinogen II 8102: + Anti-h Pepsinogen II 8103: +
--------------------------	--





Legal disclaimer

Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.