

Anti-h Copeptin 4804 SPTN-5

Product overview

Catalog number	100649
Specificity	Antibody recognizes human copeptin
Description	Monoclonal mouse antibody, cultured <i>in vitro</i> under conditions free from animal-derived components.
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9 % NaCl, 0.095 % NaN ₃ as a preservative
Shelf life and storage	Unspecified, storage at 2–8 °C
Subclass	IgG ₁
Analyte description	Copeptin is a 39-amino acid glycopeptide, cleaved from the C-terminus of pre-provasopressin (pre-proAVP). It has been suggested as a biomarker in diagnosis and prognosis of several diseases, such as acute myocardial infarction, heart failure, hyponatremia, and sepsis.

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	7.2–7.8
Purity	≥ 95 %

Kinetic parameters

Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-



Legal disclaimer

Cross-reactivities N/D

Epitope Epitope is located within the sequence QLAGAPEPFEPAPQPDAY in the C-terminal part of copeptin.

Pair recommendations

		DETECTION			
		4801	4802	4804	4806
CAPTURE	4801	-	-	+	+
	4802	-	-	+	+
	4804	+	+	-	-
	4806	+	+	-	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA

Antigens tested N/D

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	Reduced immunoreactivity
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 3 days	OK
+45 °C, 7 days	Reduced immunoreactivity

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -



Legal disclaimer