Medix Biochemica

Product specification
ANTIBODY
2026-01-09

Anti-h SCC 11801 SPTN-5

Product overview

Catalog number	100876
Specificity	Antibody recognizes human squamous cell carcinoma (SCC) antigen
Description	Monoclonal mouse antibody, cultured *in vitro* under conditions free from animal-derived components.
Product buffer solution	50 mM Na-citrate, pH 6.0, 0.9% NaCl, 0.095% NaN_3 as a preservative
Shelf life and storage	36 months from manufacturing at 2–8 °C
Subclass	IgG_1
Analyte description	Squamous cell carcinoma (SCC) antigen is a biomarker of squamous cell carcinomas found in the lung, uterine cervix, esophagus, head, neck, anal canal, and skin. In clinical applications SCC antigen levels are measured from serum or plasma to monitor recurrence of the tumor and response to the therapy. This marker is not recommended for screening, since SCC antigen levels may also increase in non-malignant diseases including severe inflammatory diseases, psoriasis, and renal failure.

Parameters tested on each lot

Product appearance	Liquid, may turn slightly opaque during storage
Product concentration	5.0 mg/ml (+/- 10 %)
Immunoreactivity	80–120 % compared to the reference sample in an FIA test
IEF Profile	6.5–7.3
Purity	≥ 95 %

Kinetic parameters

Association rate constant	Not Determined (N/D)
Dissociation rate constant	N/D
Affinity constant	N/D
Determination method	-
Determination antigen	-

Oy Medix Biochemica Ab
Klovinpellontie 3,
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI14631532

Legal disclaimer
Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.

Cross-reactivities N/D

Epitope N/D

Pair recommendations

		DETECTION			
		11801	11802	11803	11804
CAPTURE	11801	-	-	+	+
	11802	-	-	+	+
	11803	+	+	-	+
	11804	+	+	+	-

Please note that pair recommendations are based on results obtained by our laboratory. Equally good results may be obtained using other pairs and therefore these recommendations are only indicative.

Platforms tested FIA, CLIA

Antigens tested N/D

Product stability

TEMPERATURE, TIME	RESULT
-70 °C, 21 days	OK
-20 °C, 21 days	OK
+4 °C, 21 days	OK
+35 °C, 21 days	OK
+45 °C, 7 days	OK

Stability testing is performed in the product buffer to see whether different temperatures affect the antigen binding, charge or composition of the antibody. Please note that the shelf life given on the first page is based on real time stability testing at 2–8 °C in the product buffer.

Miscellaneous -

References -

Oy Medix Biochemica Ab
Klovinpellontie 3,
FI-02180 Espoo, Finland

medix@medixbiochemica.com
www.medixbiochemica.com
VAT reg. no. FI14631532

Legal disclaimer
Medix Biochemica products meet their specifications if transported, stored, and used according to the instructions. Medix Biochemica's products may not be used or reproduced without Medix Biochemica's written permission.